

Original Research

Application of Functional Tests in the Prevention from Musculoskeletal Injuries in Young Classical Style Wrestlers

Małgorzata Wójcik^{1,*}, Idzi Siatkowski², Piotr Żurek³¹Institute of Health Sciences, Hipolit Cegielski State University of Applied Sciences, 62-200 Gniezno, Poland²Department of Mathematical and Statistical Methods, Poznan University of Life Science, 60-637 Poznan, Poland³Faculty of Sport Sciences in Gorzow Wielkopolski, Poznan University of Physical Education, 61-871 Poznan, Poland*Correspondence: malgo_wojcik@interia.pl (Małgorzata Wójcik)

Submitted: 29 December 2021 Revised: 19 January 2022 Accepted: 5 February 2022 Published: 13 May 2022

Abstract

Background: Wrestling is a contact sport with a high risk of injury occurrence caused by typical motor system dysfunctions. In wrestling training a high value is put on the development of power and muscle strength, however, a program aimed at improving motor control is not introduced. **Methods:** The study involved 25 men wrestlers from School of Sporting Excellence in Radom Poland. To identify musculoskeletal system weak links low threshold Performance Matrix Tests were used. An analysis of weak links occurrence in biokinematics chain was done using ANalysis Of Variance (ANOVA), location and direction of weak links occurrence was identified. **Results:** The low threshold tests provide the information that tested athletes ($n = 25$) have weak links in musculoskeletal system. Players training experience has significant influence on their occurrence $Pr (>F) 0.01809$. The values $p_3 = Pr (>F) 0.03215$ and $p_5 = Pr (>F) 0.04042$ reflect significant correlation with occurrence of weak links number in various places in wrestlers musculoskeletal system. The results $Pr (>F)$ for training frequency, wrestlers age and sports level indicate no significant effect of these characteristics on weak links location in musculoskeletal system. **Conclusions:** Obtained results lead to the following conclusions: (1) training loads in wrestling cause weak links occurrence in musculoskeletal system among juniors; (2) wrestlers training in junior category significantly affects weak links occurrence in chain of musculoskeletal system in different locations; (3) wrestlers training frequency does not have a significant impact on weak links location estimated by low threshold tests.

Keywords: greco-roman wrestling; prevention; Performance Matrix testing; injuries; weak links; ANOVA

1. Introduction

Sports fight in wrestling is conducted under the rules of classical or free style. Wrestling is a contact sport which favors the formation of musculoskeletal dysfunction in athletes. Training and fighting in wrestling is characterized by significant risks of common musculoskeletal injuries [1–5]. In sports training in wrestling a high value is put on the development of muscular strength and power mainly through the isometric work of muscles [6], however, any actions are undertaken to apply an activity focused on improved motor control. It is forgotten that an inadequate motor control is responsible for abnormal motor pattern, which in turn reduces the effectiveness of sports training. It has been proved that there is a correlation among muscle strength, power and stability vitally needed for a proper motor control function [7,8]. Prevention of musculoskeletal dysfunction disorders is a complex process, which requires a holistic treatment of a player associated with the close cooperation with coaches, physiotherapists and doctors. The positive effect of a player's holistic treatment is motor control improving, which can give the results in lengthening the time career of a player and can reduce treatment costs. Because of the problem's importance we undertook the attempt to evaluate wrestler's musculoskeletal system using standardized Performance Matrix Tests. To assess the ef-

fectiveness of the operation of the musculoskeletal system, we proposed using tests which do not incur high financial costs and which can be carried out in any location where the athletes might be, either during normal training or just before competition. Functional evaluation allows us to introduce a therapeutic program that reduces the number of weak links and thus the probability of injury occurrence. The use of such a program in a wrestlers training cycle should enable more effective impact on the improvement of motor control and the functioning of the musculoskeletal system.

Assessing the musculoskeletal system using Performance Matrix Tests has hitherto been carried out on athletes in a range of disciplines: fencers, hockey players, canoeists and rowers, as well as among amateur athletes, such as university sports students and pupils at ballet school [9–13]. Some athletes, on identifying weak links in the musculoskeletal system, segmental stabilization training was undertaken which significantly reduced them [11]. Strong core stability reduces the likelihood of injury [14,15].

The following targets were taken into consideration: (1) to determine how the training loads in wrestling influence the presence of weak links in the musculoskeletal system; (2) to determine the relationship among the occurrence of weak links in the musculoskeletal system, training duration and training frequency in junior wrestlers in the classical style. The following research hypotheses were for-



mulated: (1) training loads in wrestlers cause the presence of weak links in the musculoskeletal system; (2) the occurrence of weak links in the musculoskeletal system depend to varying degrees on training experience, athlete's age, sports classes and frequency of training; (3) training experience, athlete's age, sports level and frequency of training do not affect the actual location of the weak links.

2. Methods

2.1 Participants

The study involved 25 young men, highly skilled wrestlers from School of Sporting Excellence in Radom Poland: age $16.7 \text{ years} \pm 1.2 \text{ years}$ (mean \pm standard deviation), body weight $77.7 \text{ kg} \pm 18.7 \text{ kg}$, body height $175.2 \text{ cm} \pm 14.1 \text{ cm}$. The average training duration of the players was $6.4 \text{ years} \pm 2.4 \text{ years}$. Among the respondents were members of the Polish Junior National Team and Polish Junior Championship medalists. Wrestlers participation in on an average in $5.4 \text{ units} \pm 0.5 \text{ units}$ of training weekly. Test subjects were volunteers, athletes preparing for the most important competition of that season—greco-roman world wrestling championships. The test subjects did not suffer injury. Injured athletes were excluded from the tests, as this could have negatively affected the data generated.

2.2 Testing Procedures

To evaluate the presence of weak links, low threshold Performance Matrix Tests (Fig. 1), which are subject to copyright, were used [16,17].

By analyzing and identifying the motion performed by a player, the weakest links of the musculoskeletal system were found. Tests were always assessed by the same investigator. The players received an explanation and shown how to execute the test. After the test, the results were inscribed in to the sheet. Five low threshold activity tests were carried out to evaluate wrestlers motor control: test 1—standing control on slightly bent leg, test 2—spine dissociation, test 3—control of the shoulder joint in the standing, test 4—limbs control with bent knee joints, test 5—limbs control in lean kneeling. The tests were carried out by an experienced physiotherapist, specially trained in conducting and assessing the results of such tests. The athletes reported no problems with carrying out the tests. Scoring two or more points on an individual test indicated a high risk of a weak point occurring in the musculoskeletal system.

2.3 Statistical Analysis

In the statistical study of the results we used one-way ANOVA (Analysis Of Variance) and multi-way ANOVA. The location of weak links is marked as: p1—cervical spine, p2—thoracic spine, p3—brachiocephalic shoulder joint, p4—shoulder joint, p5—lumbar spine, p6 - knee joint. The results were statistically analyzed using the statistical package R [18].

3. Results

Weak links in the musculoskeletal system were found in all athletes ($n = 25$) (Fig. 2). Their presence was noted at six levels: cervical, thoracic and lumbar spine, in brachiocephalic shoulder joint, the shoulder joint and knee joint. There were no weak links at the hip. Among all respondents in the cervical spine weak links were observed in the direction of bending in two players, in the thoracic spine weak link appeared in three players in the direction of bending, and in the direction of rotation in nine wrestlers. However, in the lumbar spine weak links have been observed in the direction of rotation in up to 24 respondents, among ten wrestlers in the direction of flexion and lateral flexion. In two wrestlers we observed lack of scapula control in brachiocephalic shoulder joint. In the shoulder joint weak links appeared in the direction of front slip in 19 respondents and in 10 competitors in the direction of rotation. At the level of knee weak link occurred in the direction of rotation in five respondents.

We verified the dependence between training experience, age of players, level of sport and the number of training in week micro cycle and the occurrence of weak links in the low threshold tests using ANOVA for multifactor experiments. It must be mentioned that analysis of the value $Pr (>F)$ has not shown a significant effect of wrestlers age, their level of sport and the number of training units on the presence of weak links in low threshold activity tests. While the wrestlers training experience has a significant influence on the occurrence of weak links in the musculoskeletal system $Pr (>F) 0.01809$ (Table 1).

Another ANOVA variation analysis was conducted to test hypotheses about the impact of training experience on the location of weak links. The values obtained for the location of p3 (brachiocephalic shoulder joint) $Pr (>F) 0.03215$ and p5 (lumbar spine) $Pr (>F) 0.04042$ entitle us to apply for their significant influence on the occurrence of weak links in various places of musculoskeletal system in wrestlers. It was also found that, the value $Pr (>F)$ p1 location (cervical spine), p2 (thoracic spine), p4 (shoulder joint) and p6 (knee joint) did not significantly influence the presence of weak links (Table 2).

The next step of statistical analysis was to check frequency significance of wrestling training on the location of weak links in low threshold tests. The obtained value $Pr (>F)$ indicates no significant training frequency effect on location of the players weak links (Table 3).

Another ANOVA variation analysis was conducted to verify the significance of the dependence between the age of athletes and the location of weak links occurrence (Table 4). The calculated value $Pr (>F)$ indicates no significant effect of wrestlers age on weak links' location. However, value $Pr (>F)$ for the level of p1 0.0983 and $Pr (>F) 0.0819$ for level p2, were close to the limit what proves the existence of a significant effect of players' age on weak links occurrence (Table 4).

Table 1. ANOVA effect of training experience, age of players, level of sport and the number of week micro cycle training on the presence of weak links in low threshold tests.

Locations	Df	SumSq	MeanSq	Fvalue	Pr (>F)
Training experience	1	0.26265	0.26265	6.629	0.01809*
Age	1	0.10225	0.10225	2.5807	0.12385
Sports level	1	0.02631	0.02631	0.6639	0.42477
Number of training sessions per week	1	0.00003	2.5E-05	0.0006	0.98016
Residuals	20	0.79244	0.03962		

Signif. codes of p -values for $P(>F)$: 0 '****' 0.001 '***' 0.01 '**' 0.05 '.' 0.1 ' ' 1.



Fig. 1. Example of low threshold Performance Matrix Test.

Subsequently, ANOVA was conducted to assess the impact of sports level on weak links location (Table 5). The obtained results of $Pr (>F)$ show no significant impact of sports level of athletes on the location of weak links in musculoskeletal system.

Table 2. ANOVA effect of wrestlers training experience on the location of weak links in low threshold tests.

Locations	Df	SumSq	MeanSq	Fvalue	Pr (>F)
p1	1	0.043	0.0426	0.0107	0.91892
p2	1	5.691	5.6915	1.4236	0.2483
p3	1	21.558	21.5584	5.3924	0.03215*
p4	1	12.042	12.0415	3.0119	0.09974
p5	1	19.499	19.499	4.8773	0.04042*
p6	1	4.964	4.9641	1.2417	0.27981
Residuals	18	71.963	3.9979		

Signif. codes of p -values for $P(>F)$: 0 '****' 0.001 '***' 0.01 '**' 0.05 '.' 0.1 ' ' 1.

Table 3. ANOVA effect of wrestling training frequency on the location of weak links in low threshold tests.

Locations	Df	SumSq	MeanSq	Fvalue	Pr (>F)
p1	1	0.2817	0.28174	1.2161	0.2846
p2	1	0.1449	0.14493	0.6256	0.4393
p3	1	0.3313	0.3313	1.4301	0.2473
p4	1	0.636	0.63598	2.7452	0.1149
p5	1	0.0009	0.00086	0.0037	0.9521
p6	1	0.1952	0.19518	0.8425	0.3708
Residuals	18	4.17	0.23167		

Signif. codes of p -values for $P(>F)$: 0 '****' 0.001 '***' 0.01 '**' 0.05 '.' 0.1 ' ' 1.

4. Discussion

The study has confirmed the first research hypothesis that the training loads in wrestling cause the occurrence of weak links (set by low threshold test) in the musculoskeletal system (Fig. 2). Particularly important in the players evaluation is the ability to indicate the place in musculoskeletal system, in which a weak link can occur. This knowledge helps to improve a prevention from a player's injury. Such detailed and comprehensive analysis of wrestlers' musculoskeletal system also allows us to apply a precise treatment program, which is an important addition to sports training.




Fig. 2. Boxplot of weak links numbers of tested wrestlers (n = 25).

Table 4. ANOVA effect of wrestlers age on the location of weak links in low threshold tests.

Locations	Df	SumSq	MeanSq	Fvalue	Pr (>F)
p1	1	4.0209	4.0209	3.0391	0.0983
p2	1	4.491	4.491	3.3944	0.0819
p3	1	2.0392	2.0392	1.5413	0.2304
p4	1	0.4571	0.4571	0.3455	0.5639
p5	1	0.0108	0.0108	0.0081	0.9292
p6	1	0.9261	0.9261	0.6999	0.4138
Residuals	18	23.815	1.3231		

Signif. codes of p -values for $P(>F)$: 0 '****' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1.

Table 5. ANOVA effect of wrestlers sports level on location of weak links in low threshold tests.

Locations	Df	SumSq	MeanSq	Fvalue	Pr (>F)
p1	1	0.3835	0.38348	0.5794	0.4564
p2	1	0.0195	0.01948	0.0294	0.8657
p3	1	0.4583	0.45834	0.6925	0.4162
p4	1	1.0392	1.03916	1.5701	0.2262
p5	1	0.0008	0.00084	0.0013	0.972
p6	1	0.0254	0.02541	0.0384	0.8469
Residuals	18	11.9133	0.66185		

Signif. codes of p -values for $P(>F)$: 0 '****' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1.

Research hypotheses have been verified which suggest that the presence of weak links in the musculoskeletal system does not depend on wrestlers age, their performance level and frequency of participation in training. However, it has been found that training experience significantly affects the occurrence of weak links $Pr(>F)$ 0.01809 (Table 1).

Also we examined whether the training experience, wrestlers age, sports level and frequency of workouts do not have a significant effect on the location of the weak links. On the basis of the value $Pr(>F)$ presented in Table 2 we can assure that for the location p1 (cervical spine), p2 (thoracic spine), p4 (shoulder) and p6 (knee) (Table 2) training duration has no an significant effect on the occurrence of weak links in musculoskeletal system. However, for location p3 (brachiocephalic shoulder joint) and p5 (lumbar spine) has a significant impact because for p3 = $Pr(>F)$ and 0.03215 and for p5 = $Pr(>F)$ 0.04042 (Table 2).

The obtained results give us valuable information pointing at the necessity of introducing a preventive action for wrestlers with training experience $6.36 \text{ years} \pm 2.38 \text{ years}$ which would block the occurrence of weak links in the musculoskeletal system among the respondents.

Frequency of training determined on the base of a number of training units in the micro cycle, shows no significant effect on the location of the weak links in musculoskeletal system (Table 3). Also the value $Pr(>F)$ for age and sports level does not significantly influence the location of the weak links (Tables 4,5).

Performance Matrix Tests could serve as a reliable research tool, for example as part of selection tests prior to athletes commencing a training period [17,19]. It has been shown that problems in the neuromuscular system in the core is a contributing factor to the occurrence of spinal sports injuries, as well as in specific sections of the kinetic chain. The strategy for preventing injury relates to proper control, and so to the correct neuromuscular transmission [20].

Martial arts have a high risk of injuries [4,5,21,22]. The issue related to incidence of sports injuries among wrestlers is relatively well known. The most frequently mentioned locations of injuries in musculoskeletal system in wrestlers are knee joints, shoulder joints, head and face, lumbar spine, ankles, cervical spine and muscle strain [4,5,23–30]. Muscle rupture or strain is also a cause of a substantial percentage of injuries among players in different sports disciplines [31].

In martial arts much effort is put into working on the increase in muscle power and strength. More frequently applied kind of training in sport, according to Core Stability concept, concerns working on shaping local and global trunk muscles, upper and lower limbs, in symmetrical and asymmetrical movements, combined with trunk muscles' activity [17].

Our study has provided the information that places in musculoskeletal system with a weaker motor control in tested players are brachiocephalic shoulder joint and lumbar spine. More frequent injuries of elbow and knee joints, head, face and nose are among the competitors practicing the classical wrestling style [32]. It should be mentioned that not enough sufficient attention is paid to the prevention from injuries among wrestlers. Most coaches working with players focus their attention on the development and improving sports technique [33]. Certainly, greater coaches awareness on the use of injury prevention and the introduction of these activities in training practice would contribute to reducing the occurrence of injuries in wrestlers [34].

Changes in the strategies and techniques of fighting could provide a way of avoiding or limiting the occurrence of injuries [35]. It is extremely important to develop a training model (including tactical-technical elements) aimed at achieving a high sporting level while reducing the occurrence of injuries in athletes [36]. The multifaceted nature of training wrestling as an Olympic discipline should be rooted in predicting possible injuries and their prevention. Martial arts, including wrestling, have a high incidence of injury. Introducing tests which may quickly and easily assess the existence of a weak link in the musculoskeletal system could aid athletes in their preparations for competition (avoiding injury) and be one of the elements influencing the attainment of a high sporting level.

5. Conclusions

Training loads in the classic wrestling style cause the occurrence of weak links in musculoskeletal chain among juniors. Wrestlers training duration in the junior category significantly affects the occurrence of weak links in the musculoskeletal chain in different locations. Wrestlers training frequency do not have a significant impact on the location of the weak links in musculoskeletal system evaluated by using low threshold tests.

Author Contributions

Conceptualization—MW, IS and PŻ; data curation—MW; formal analysis—MW, IS and PŻ; statistical analysis, software—IS; investigation—IS; methodology—MW and PŻ; project administration—MW; supervision—MW, IS and PŻ; validation—MW, IS and PŻ; visualization—MW and PŻ; writing - original draft—MW, IS and PŻ; writing - review & editing—MW, IS and PŻ. All authors have read and agreed to the published version of the manuscript.

Ethics Approval and Consent to Participate

All procedures performed in studies involving human participants were in accordance with the ethical standards of institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all individual participants included in the study. The study protocol was approved by the Bioethics Committee at the Medical University of Poznan, Poland (permit No. 509/10).

Acknowledgment

Thank numerous individuals participated in this study.

Funding

This research received no external funding.

Conflict of Interest

The authors declare no conflict of interest.

References

- [1] Yard EE, Comstock RD. A comparison of pediatric freestyle and Greco-Roman wrestling injuries sustained during a 2006 us national tournament. *Scandinavian Journal of Medicine and Science in Sports*. 2008; 18: 491–497.
- [2] Barosso BG, da Silva JMA, da Costa Garcia A, de Oliveira Ramos NC, Martinelli MO, Resende VR, *et al.* Musculoskeletal injuries in wrestling athletes. *Acta Ortopedica Brasileira*. 2011; 19: 98–101.
- [3] Stanev S, Dimitrova E. Training risk factors associated with wrestling injury. *British Journal of Sports Medicine*. 2011; 45: 80–81.
- [4] Park KJ, Lee JH, Kim HC. Injuries in male and female elite Korean wrestling athletes: a 10-year epidemiological study. *British Journal of Sports Medicine*. 2019; 53: 430–435.
- [5] Kim JC, Park KJ. Injuries and rapid weight loss in elite Korean wrestlers: an epidemiological study. *The Physician and Sportsmedicine*. 2021; 49: 308–315.
- [6] McGuigan MR, Winchester JB, Erickson T. The importance of isometric maximum strength in college wrestlers. *Journal of Sports Science and Medicine*. 2006; CSSI: 108–111.
- [7] Comerford MJ, Mottram SL. Movement and stability dysfunction – contemporary developments. *Manual Therapy*. 2001; 6: 15–26.
- [8] McGill SM. *Low Back Disorders: Evidence - based prevention and rehabilitation* (pp. 109). 2nd edn. Champaign Human Kinetics Publishers: Champaign. 2007.
- [9] Mischiati C, Comerford M, Gosford E, Swart J, Ewings S, Botha N, *et al.* Intra and Inter-Rater Reliability of Screening for Movement Impairments: Movement Control Tests from The Foundation Matrix. *Journal of Sports Science and Medicine*. 2015; 14: 427–440.
- [10] Wójcik M, Siatkowski I. The Usefulness of Performance Matrix Tests in Locomotor System Evaluation of Girls Attending a Ballet School — Preliminary Observation. *Journal of Physical Therapy Science*. 2014; 26: 41–44.
- [11] Wójcik M, Siatkowski I, Bodnar-Nanuś A. The influence of segmental stabilization training upon the reduction of motor system weak connections in hockey players. *Chirurgia Narządów Ruchu i Ortopedia Polska*. 2011; 76: 145–150.

- [12] Wójcik M, Siatkowski I. Occurrence of Weak Links of Biokinematics Chain and Pain Feeling in the Lumbar Spine in Kayakers and Female Rowers – Preliminary Observation. *Rheumatology*. 2011; 2161–1149.
- [13] Wójcik M, Siatkowski I, Nowakowski A, Witkowski M. Performance Matrix tests usage for diagnosing weak links of musculoskeletal system occurred in fencers. *Chirurgia Narządów Ruchu i Ortopedia Polska*. 2011; 76: 41–46.
- [14] Comerford M. Core stability: priorities in rehabilitation of the athlete. *SportEx Medicine*. 2004; 22: 15–22.
- [15] Comerford M. Screening to identify injury and performance risk: movement control testing—the missing piece of the puzzle. *SportEx Medicine*. 2006; 29: 21–26.
- [16] Hodges PW, Moseley GL. Pain and motor control of the lumbopelvic region: effect and possible mechanisms. *Journal of Electromyography and Kinesiology*. 2003; 4: 361–370.
- [17] Mottram S, Comerford M. A new perspective on risk assessment. *Physical Therapy in Sport*. 2008; 9: 40–51.
- [18] R Development Core Team. A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria. 2009. Available at: <http://www.r-project.org> (Accessed: 10 September 2021).
- [19] McNeill W. The Double Knee Swing Test - a practical example of the Performance Matrix Movement Screen. *Journal of Bodywork and Movement Therapies*. 2014; 18: 477–481.
- [20] Zazulak B, Cholewicki J, Reeves NP. Neuromuscular control of trunk stability: clinical implications for sports injury prevention. *The Journal of the American Academy of Orthopaedic Surgeons*. 2008; 16: 497–505.
- [21] Graham MR, Davies B, Baker JS. Head injuries in full contact karate competition! Is the prejudice in management minimising the required investigation? *Journal of Sports Science and Medicine*. 2007; 6: 62–64.
- [22] Halabchi F, Ziaee V, Lotfian S. Injury profile in women shotokan karate championships in iran (2004-2005). *Journal of Sports Science and Medicine*. 2007; 6: 52–57.
- [23] Boden BP, Lin W, Young M, Mueller FO. Catastrophic injuries in wrestlers. *The American Journal of Sports Medicine*. 2002; 30: 791–795.
- [24] Lightfoot AJ, McKinley T, Doyle M, Amendola A. ACL Tears in Collegiate Wrestlers. *The Iowa Orthopaedic Journal*. 2005; 25: 145–148.
- [25] Agel J, Ransone J, Dick R, Oppliger R, Marshall SW. Descriptive epidemiology of collegiate men's wrestling injuries: National Collegiate Athletic Association Injury Surveillance System, 1988–1989 through 2003–2004. *Journal of Athletic Training*. 2007; 42: 303–310.
- [26] Michael JW, Müller L, Schikora N, Eysel P, König DP. Injuries to the upper limbs in competitive wrestlers. *Sportverletzung Sportschaden*. 2008; 22: 89–92. (In German)
- [27] Bonza JE, Fields SK, Yard EE, Dawn Comstock R. Shoulder Injuries among United States High School Athletes during the 2005–2006 and 2006–2007 School Years. *Journal of Athletic Training*. 2009; 44: 76–83.
- [28] Myers RJ, Linakis SW, Mello MJ, Linakis JG. Competitive Wrestling-related Injuries in School Aged Athletes in U.S. Emergency Departments. *The Western Journal of Emergency Medicine*. 2010; 11: 442–449.
- [29] Aydos L. Effect of wrestling on the foot sole of elite wrestlers. *International Journal of the Physical Sciences*. 2011; 6: 3143–3154.
- [30] Thomas RE, Zamanpour K. Injuries in wrestling: systematic review. *The Physician and Sportsmedicine*. 2018; 46: 168–196.
- [31] McNally EG. *Ultrasonografia narządu ruchu* (pp. 223). 1st edn. Elsevier Urban&Partner: Wrocław. 2008. (In Polish)
- [32] Yard EE, Collins CL, Dick RW, Comstock RD. An epidemiologic comparison of high school and college wrestling injuries. *The American Journal of Sports Medicine*. 2008; 36: 57–64.
- [33] Sadowski J, Gierczuk D. Effectiveness of coordination training in technical preparation of freestyle wrestlers at various levels of sports advancement. *Archives of Budo*. 2010; 6: 143–148
- [34] Boostani MH, Boostani MA, Rezaei AM, Khatamsaz S. The study of sport injuries in the national team karate of Iran (kumite field). *Journal of Martial Arts Anthropology*. 2011; 11: 27–34.
- [35] Miarka B. Demandas técnicas e fisiológicas de combates da luta olímpica. *Revista de Artes Marciales Asiáticas*. 2016; 11: 18–31.
- [36] Vasilescu F, Leonte N, Tudorancea SD. Contributions regarding the development and experimentation of a specific physical training model for greco-roman junior wrestlers. *Revista Românească pentru Educație Multidimensională*. 2021; 13: 442–456.